

Childhood brain tumors develop early in highly specialized nerve cells

Medulloblastomas, childhood brain tumors in children, are thought to develop between the first trimester of pregnancy and the end of the first year of life. Researchers at the Hopp Children's Cancer Center Heidelberg (KiTZ), the German Cancer Research Center (DKFZ) and Heidelberg University Hospital (UKHD) have now published their findings in the journal *Nature*. They analyzed the genetic changes of each individual cancer cell in tumor samples in order to reconstruct which genetic changes occur first during tumor development and when.

Medulloblastoma is one of the most common malignant tumors of the central nervous system (CNS) in children and adolescents. It develops in the area of the cerebellum, which is responsible for movement coordination, among other things. Medulloblastomas enlarge rapidly, often grow into surrounding tissue and can also form metastases. The wide variety of this tumor group also makes it difficult to find the right treatment.

A team of researchers at the Hopp Children's Cancer Center Heidelberg (KiTZ), the German Cancer Research Center (DKFZ) and Heidelberg University Hospital (UKHD) has now investigated exactly where and when the most aggressive subgroups of these tumors develop in so-called single cell analyses. "We have genetically characterized the many thousands of individual cells from the tumor samples of numerous young patients with medulloblastoma," explains Konstantin Okonechnikov, first author of the study from KiTZ and DKFZ. "This technology allows us to obtain a high-resolution picture of the genetic make-up within a tumor, so that we can reconstruct its developmental history fairly accurately."

Accordingly, there are "early" and "late" genetic changes within a tumor. Based on the distribution of the genetic changes across the various tumor cell clones, the scientists were able to reconstruct the development of the tumor. The study shows that the particularly aggressive medulloblastomas of subgroups three and four probably develop between the first trimester of pregnancy and the end of the first year of life. The origin of the tumor development are the precursor cells of highly specialized nerve cells, the so-called unipolar brush cells of the cerebellum, which also develop between the first trimester of pregnancy and the end of the first year of life. Large rearrangements of entire chromosomes or chromosome arms take place in these cells, presumably initially by chance, as the present results show: The cells lose or gain certain chromosomes.

"We assume that the early loss or gain of certain chromosomes is the first step in tumor development and that these occur many years before the clinical symptoms appear," says Lena Kutscher from KiTZ and DKFZ, who led the study together with Stefan Pfister, Director at KiTZ, Head of Department at DKFZ and paediatric oncologist at Heidelberg University Hospital (UKHD).

Only the later tumor cells carry the already known duplications or alterations of the cancer genes MYC, MYCN or PRDM6 that are typical for these types of medulloblastoma. "We therefore assume that these cancer genes are responsible for the progressive tumor growth and also for the metastasis and therapy resistance that occurs, but not for the development of the tumor," says Lena Kutscher.

Lena Kutscher explains what these results could mean for diagnosis and treatment: "If we succeed in developing sufficiently sensitive methods in the future to detect these early changes, for example as DNA fragments in the blood, this could form the basis for possible early detection in newborns and infants."

Publication

K. Okonechnikov *et al.*: Oncogene aberrations drive medulloblastoma progression, not initiation. In: *Nature* (Online Publikation 7. Mai 2025). DOI: 10.1038/s41586-025-08973-5

08-May-2025

Source: German Cancer Research Center (DKFZ)

Further information

- ▶ German Cancer Research Center (DKFZ)